arXiv:1002.0588v22 [math.CO] 10 Oct 2011

The distance-3 graph of the Biggs-Smith graph

Italo J. Dejter
University of Puerto Rico
Rio Piedras, PR 00936-8377
ijdejter@uprrp.edu

Abstract

The distance-3 graph $\mathcal{S}^3$ of the Biggs-Smith graph $\mathcal{S}$ is shown to be: **(a)** a connected edge-disjoint union of 102 tetrahedra (copies of K_4) and as such the K_4-ultrahomogeneous Menger graph of a self-dual (102_4)-configuration; **(b)** a union of 102 cuboctahedra, (copies of $L(Q_3)$), with no two such cuboctahedra having a common chordless 4-cycle; **(c)** not a line graph. Moreover, $\mathcal{S}^3$ is shown to have a $\mathcal{C}$-ultrahomogeneous property for $\mathcal{C} = \{K_4\} \cup \{L(Q_3)\}$ restricted to preserving a specific edge partition of $L(Q_3)$ into 2-paths, with each triangle (resp. edge) shared by two copies of $L(Q_3)$ plus one of K_4 (resp. four copies of $L(Q_4)$). Both the distance-2 and distance-4 graphs, $\mathcal{S}^2$ and $\mathcal{S}^4$, of $\mathcal{S}$ appear in the context associated with the above mentioned edge partition. This takes us to ask whether there any non-line-graphical connected K_4-ultrahomogeneous Menger graphs of self-dual (n_4)-configurations that are edge-disjoint unions of at least two copies of K_4 for positive integers $n \notin \{42, 102\}$.

1 Introduction

Given a (di)graph Γ and a positive integer $k \leq$ diameter(Γ), the *distance-k graph* Γ^k of Γ has $V(\Gamma^k) = V(\Gamma)$ and an arc in Γ^k from a vertex u to a vertex $v \neq u$ whenever there is a shortest k-arc of length k in Γ from u to v. A k-arc in a graph is a sequence of vertices $(v_0, \dots, v_k)$ such that consecutive vertices are adjacent and $v_{i-1} \neq v_{i+1}$ when $0 < i < k$ [9]. A k-arc can also be interpreted as a directed walk of length k in which consecutive edges are distinct [11].

Ultrahomogeneous (or UH) graphs were introduced and treated in [3, 8, 10, 14, 15], but we deal here with the following modified concept of ultrahomogeneity. Given a collection $\mathcal{C}$ of (di)graphs closed under isomorphisms, a (di)graph G is said to be *$\mathcal{C}$-UH* if every isomorphism between two induced members of $\mathcal{C}$ in G extends to an automorphism of G. If $\mathcal{C}$ is the isomorphism class of a graph H, then G is said to be $\{H\}$-UH or H-UH. If $\mathcal{C}$ is the isomorphism class of a graph H with an edge partition Ω into 2-paths), then G is said to be *Ω-preserving* $\{H\}$-UH or H-UH if only every Ω-preserving isomorphism between two induced copies of H extends to an automorphism of G. In [12], $\mathcal{C}$-UH graphs are defined and studied when $\mathcal{C}$ is the collection formed either by the complete graphs, or the disjoint unions of complete graphs, or the complements of those disjoint unions.

Let M be a subgraph of a graph H and let G be both an M-UH and an H-UH graph. (A particular case has $H = L(Q_3)$ with an edge partition Ω into 2-paths, as for example the $L(Q_3)$ represented in Figure 4, Section 4 below, with M equal to a 2-path in Ω). We say that G is an (Ω-*preserving*) $\{H\}_M$-*UH graph* if, for each induced copy H_0 of H in G containing an induced copy M_0 of M, there exists exactly one induced copy $H_1 \neq H_0$ of H in G with $V(H_0) \cap V(H_1) = V(M_0)$ and $E(H_0) \cap E(H_1) = E(M_0)$. These vertex and edge conditions can be condensed as $H_0 \cap H_1 = M_0$. We say that such a G is *tightly fastened*. This is generalized by saying that an $\{H\}_M$-UH graph G is an ℓ-*fastened* $\{H\}_M$-UH graph if given an induced copy H_0 of H in G containing an induced copy M_0 of M, then there exist exactly ℓ induced copies $H_i \neq H_0$ of H in G such that $H_i \cap H_0 \supseteq M_0$, for each $i = 1, 2, \ldots, \ell$, with $H_1 \cap H_0 = M_0$. Observe that in these definitions we started by assuming that G is M-UH and H-UH.

A transformation of distance-transitive graphs into $\mathcal{C}$-UH graphs that goes from the Coxeter graph on 28 vertices into the Klein graph on 56 vertices [7] is applied below to the Biggs-Smith graph $\mathcal{S}$ [2, 4] seen as a $\{C_9\}_{P_4}$-UH graph. This allows to show that the graph obtained by the said transformation, namely the distance-3 graph $\mathcal{S}^3$ of $\mathcal{S}$, is a connected edge-disjoint union of 102 tetrahedra (copies of K_4) as well as a union of 102 cuboctahedra, (copies of $L(Q_3)$) with no two such cuboctahedra having a common 4-hole (chordless 4-cycle). In addition, $\mathcal{S}^3$ is shown to possess a $\mathcal{C}$-UH property for $\mathcal{C} = \{K_4\} \cup \{L(Q_3)\}$ restricted to preserving a specific edge partition of $L(Q_3)$ into 2-paths, with each triangle (resp. edge) of $\mathcal{S}^3$ shared by two copies of $L(Q_3)$ plus one of K_4 (resp. four copies of $L(Q_3)$) exactly. Moreover, $\mathcal{S}^3$ is seen to be the Menger graph [5] of a self-dual (102_4)-configuration, but not a line graph. Both $\mathcal{S}^2$ and $\mathcal{S}^4$ are shown to be related to the above mentioned edge partition.

Thus, the connected edge-disjoint union $\mathcal{X}$ of 42 copies of K_4 in [6], forming a K_4-UH graph which is the Menger graph of a (42_4)-configuration but is not a line graph, is not the only case of Menger graph of a self-dual (n_4)-configuration which is connected edge-disjoint union of copies of K_4 and K_4-UH but not a line graph: $\mathcal{S}^3$ provides another such graph. This allows to raise the following question.

Question 1 *Are there any non-line-graphical connected K_4-UH Menger graphs of self-dual (n_4)-configurations that are edge-disjoint unions of at least two copies of K_4, for positive integers $n \notin \{42, 102\}$?*

We recall that the line graph of the d-cube, is K_d-UH, where $3 \leq d \in \mathbf{Z}$ [6]. Other considerations related to Question 1 can be found in Section 5.

2 Preliminaries

We recall that the Biggs-Smith graph $\mathcal{S}$ has order $n = 102$, diameter $d = 7$, girth $g = 9$ and automorphism group $\mathcal{A} = PSL(2, 17)$ which is also shown to be that of the distance-3 graph $\mathcal{S}^3$ of $\mathcal{S}$ in Theorem 7. Let k be the largest integer s such that $\mathcal{S}$ is s-arc transitive. Then $k = 4$. Finally, we also recall that the number η of 9-cycles of $\mathcal{S}$ is $\eta = 136$.

Given a finite graph H and a subgraph M of H with $|V(H)| > 3$, a graph Γ is *strongly zipped* (or *SZ*) $\{H\}_M$*-UH* if for each sequence of connected subgraphs $M = M_1, M_2 \dots, M_t \equiv K_2$ such that M_{i+1} is obtained from M_i by the deletion of a vertex, for $i = 1, \dots, t-1$, it holds that G is a $(2^i - 1)$-fastened $\{H\}_{M_i}$-UH graph, for $i = 1, \dots, t$.

Let P_k be a $(k-1)$-path. Let C_g be a cycle of length g. Theorem 1 below asserts that $\mathcal{S}$ is SZ $\{C_g\}_{P_k}$-UH, namely SZ $\{C_9\}_{P_4}$-UH. Also, $\mathcal{S}^3$ is SZ $\{H\}_M$-UH (Theorems 6-7), where $H = L(Q_3)$ and M is a triangle, but some additional concepts are needed: A graph G is rK_s*-frequent* if every edge e of G is intersection in G of exactly r induced copies of K_s, and these copies have only e and its ends in common. (For example, K_4 is $2K_3$-frequent, and $L(Q_3)$ is $1K_3$-frequent). A graph G is $\{H_2, H_1\}_{K_3}$*-UH*, where H_i is iK_3-frequent, $(i = 1, 2)$, if: **(a)** G is an H_2-UH graph and an edge-disjoint union of induced copies of H_2; **(b)** there exists an edge partition Ω of H_1 into 2-paths, and G is Ω-preserving SZ $\{H_1\}_{K_3}$-UH; **(c)** each induced copy of H_2 in G has each of its induced copies of K_3 in common with exactly *two* induced copies of H_1 in G.

Properties of $\mathcal{S}$ needed in the rest of the paper are covered in Section 3. In Section 4, consideration of the distance-3 graphs of the 9-cycles of $\mathcal{S}$ yields a $\{K_4, L(Q_3)\}_{K_3}$-UH graph $\mathcal{Y}$ via zipping (meaning identification into an edge) of corresponding oppositely oriented arcs of the resulting 3-cycles. Theorem 7 uses Theorem 2, that establishes an explicit 1-1 correspondence between the 18 disjoint unions of 4 17-cycles each in $\mathcal{S}$ and the points of the projective line of the field $GF(17)$. Theorem 8, establishing $\mathcal{Y} = \mathcal{S}^3$, also yields two other graphs, appearing in the way the 102 copies of $L(Q_3)$ intersect, namely $\mathcal{S}^2$ and $\mathcal{S}^4$.

3 Properties of the Biggs-Smith graph $\mathcal{S}$

Let $A = (A_0, A_1, \dots, A_g)$, $D = (D_0, D_2, \dots, D_f)$, $C = (C_0, C_4, \dots, C_d)$, $F = (F_0, F_8, \dots, F_9)$ be 4 disjoint 17-cycles. Each $y = A, D, C, F$ has vertices y_i with i expressed as an heptadecimal index up to $g = 16$. We assume that i is advancing in 1,2,4,8 units mod 17, stepwise from left to right, respectively for $y = A, D, C, F$. Then the Biggs-Smith graph $\mathcal{S}$ is obtained by adding to the disjoint union $A \cup D \cup C \cup F$, for each $i \in \mathbf{Z}_{17}$, a 6-vertex tree T_i formed by the edge-disjoint union of paths $A_iB_iC_i$, $D_iE_iF_i$, and B_iE_i, where the vertices A_i, D_i, C_i, F_i are already present in the cycles A, D, C, F, respectively, and where the vertices B_i and E_i are new and introduced with the purpose of defining the tree T_i, for $0 \leq i \leq g = 16$. A corresponding Frucht diagram [13] of $\mathcal{S}$ via $\mathbf{Z}_{17}$ is depicted on the left of Figure 2 below, with nonzero Frucht notation over the loops, and null Frucht notation over the remaining edges. Now, $\mathcal{S}$ has the collection $\mathcal{C}_9$ of 9-cycles formed by:

$$\begin{array}{l|l}
S^0{=}(A_0A_1B_1C_1C_5C_9C_dC_0B_0), & W^0{=}(A_0A_1B_1E_1F_1F_9F_0E_0B_0), \\
T^0{=}(C_0C_4B_4A_4A_3A_2A_1A_0B_0), & X^0{=}(C_0C_4B_4E_4D_4D_2D_0E_0B_0), \\
U^0{=}(E_0F_0\ F_9F_1F_aF_2E_2D_2D_0,) & Y^0{=}(E_0B_0A_0A_1A_2B_2E_2D_2D_0), \\
V^0{=}(E_0D_0D_2D_4D_6D_8E_8F_8F_0), & Z^0{=}(F_0\ F_8\ E_8B_8C_8C_4C_0B_0E_0),
\end{array}$$

and those 9-cycles obtained from these, as $S^x, \ldots, Z^x$, by uniformly adding $x \in \mathbf{Z}_{17}$ mod 17 to all subindexes i of vertices y_i, so that $|\mathcal{C}_9| = 136$.

An auxiliary table presenting the form in which the 9-cycles above share the vertex sets of 3-arcs, either oppositely oriented or not, is shown below. The table details, for each one of the 9-cycles $\xi = S^0, \ldots, Z^0$, (expressed as $\xi = (\xi_0, \ldots, \xi_8)$ in the shown vertex notation), each 9-cycle η in $\{S^i, \ldots, Z^i; i = 0, \ldots, 16\} \setminus \{\xi\}$ intersecting ξ in the succeeding 3-paths $\xi_i\xi_{i+1}\xi_{i+2}\xi_{i+3}$, for $i = 0, \ldots, 8$, with additions involving i taken mod 9. Each such η in the auxiliary table has either a preceding minus sign, if the corresponding 3-arcs in ξ and η are oppositely oriented, or not, otherwise. Each shown $-\eta_j$ (resp. η_j) in the table has the subindex j indicating the equality of initial vertices $\eta_j = \xi_{i+3}$ (resp. $\eta_j = \xi_i$) of those 3-arcs, for $i = 0, \ldots, 8$:

$$\begin{array}{l|l}
S^0{:}(-T_1^e,\ T_7^1,-Z_4^1,S_4^d,\ S_3^4,-Z_3^9,\ T_0^d,-T_6^0,\ U_8^0), & W^0{:}(-U_4^0,W_2^8,W_1^9,-U_3^1,X_7^2,-X_4^b,Y_6^0,-X_8^0,X_5^9), \\
T^0{:}(\ S_6^4,-S_0^3,-Y_2^2,\ T_4^1,T_3^g,-Y_1^0,-S_7^0,\ S_1^g,\ V_8^0), & X^0{:}(-V_4^0,X_2^f,X_1^2,-V_3^4,-W_5^6,W_8^8,-Z_8^0,W_4^f,-W_7^0), \\
U^0{:}(Y_3^g,-U_6^1,Z_7^1,-W_3^g,-W_0^0,Z_0^9,-U_1^g,Y_0^0,S_8^0), & Y^0{:}(\ U_7^0,-T_5^0,-T_2^f,\ U_0^1,-Y_8^2,\ V_2^f,W_6^0,V_5^0,-Y_4^f), \\
V^0{:}(-Z_2^d,-V_6^4,Y_5^2,-X_3^d,-X_0^0,Y_7^0,-V_1^d,-Z_5^0,T_8^0), & Z^0{:}(\ U_5^8,-Z_6^8,-V_0^4,-S_5^8,-S_2^g,-V_7^0,-Z_1^9,\ U_2^g,-X_6^0),
\end{array}$$

This table is extended to all of $\mathcal{C}_9$ by adding $x \in \mathbf{Z}_{17}$ uniformly mod 17 to all superindexes.

Theorem 2 *$\mathcal{S}$ is an* SZ $\{C_9\}_{P_{i+2}}$-UH *graph, for $i = 0, 1, \ldots, k-2 = 2$. In particular, $\mathcal{S}$ is a $\{C_9\}_{P_4}$-UH graph and has exactly $2^{k-2}3n9^{-1} = 136$ 9-cycles.*

Proof. We have to see that $\mathcal{S}$ is a $(2^{i+1}-1)$-fastened $\{C_9\}_{P_{4-i}}$-UH graph, for $i = 0, 1, 2$. In fact, each $(3-i)$-path P_{4-i} of $\mathcal{S}$ is shared exactly by 2^{i+1} 9-cycles of $\mathcal{S}$, for $i = 0, 1, 2$. For $k = 4$, any edge (resp. 2-path, resp. 3-path) of G is shared by 8 (resp. 4, resp. 2) 9-cycles of G. This means that a 9-cycle C_9 of G shares a P_2 (resp. P_3, resp. P_4) with exactly other 7 (resp. 3, resp. 1) 9-cycles. The statement follows from this and the details previous to it together with a simple counting argument for the number of 9-cycles. $\square$

The reconstruction of $\mathcal{S}$ from the 17-cycles A, D, C, F will be denoted

$$\aleph(A, D, C, F) = \aleph(P^1, P^2, P^4, P^8),$$

where $P^1 = A$, $P^2 = D$, $P^4 = C$, $P^8 = F$, in order to see that it yields 8 automorphisms of $\mathcal{S}$ obtained by multiplying modulo 17 the successive exponents of P, $(= P^1 = A)$, by 2:

$$\aleph(P^1,P^2,P^4,P^8) \rightarrow \aleph(P^2,P^4,P^8,P^{-1}) \rightarrow \ldots \rightarrow \aleph(P^{-8},P^1,P^2,P^4) \rightarrow \aleph(P^1,P^2,P^4,P^8).$$

Observe that a 9-cycle x of $\mathcal{S}$ sharing exactly a 1-path x_A with A shares exactly a 2-path x_F with F, with distance 3 between x_A and x_B realized by two vertex-disjoint 3-paths in x. Note that x_A, x_F and the two 3-paths separating them are edge-disjoint. By using repeatedly this fact, it can be seen , say via a representation of $\mathcal{S}$ showing the vertices of A, D, C, F symmetrically distributed on four concentric circles, that there is another reconstruction $\aleph(A^0, D^0, C^0, F^0)$ of $\mathcal{S}$ with four disjoint 17-cycles:

$$\begin{array}{l}
A^0=(A_0A_1\;B_1C_1C_5B_5E_5F_5F_dF_4\;F_c\;E_c\;B_cC_c\;C_g\,B_g\,A_g)=\;(A^0_0,A^0_1,\ldots,A^0_g),\\
D^0=(F_0F_8\;E_8D_8D_6E_6B_6C_6C_2C_fC_b\;B_b\;E_b\;D_bD_9E_9F_9)=(D^0_0,D^0_1,\ldots,D^0_g),\\
C^0=(C_0C_4B_4A_4A_3B_3E_3D_3D_1D_gD_eE_eB_eA_eA_dB_dC_d)=\;(C^0_0,C^0_1,\ldots,C^0_g),\\
F^0=(D_0D_2E_2F_2F_aE_aB_aA_aA_9A_8A_7B_7\;E_7F_7F_fE_f\;D_f)=\;(F^0_0,F^0_1,\ldots,F^0_g),
\end{array}$$

where $A_0 = A^0_0$, $F_0 = D^0_0$, $C_0 = C^0_0$, $D_0 = F^0_0$, etc., and where the vertices A^0_i, D^0_i, C^0_i, F^0_i, for each fixed $i \in \mathbf{Z}_{17}$, are the degree-1 vertices of a 6-vertex tree T^0_i, like the tree T_i above, formed by the disjoint union of the paths $A^0_iB^0_iC^0_i$, $D^0_iE^0_iF^0_i$ and $B^0_iE^0_i$. An involution for $i \in \mathbf{Z}_{17}$ in $\aleph(A^0, D^0, C^0, F^0)$ is also present whose orbits are $\{y^0\}$, $\{y^1, y^g\}$, $\{y^2, y^f\}$, $\{y^3, y^e\}$, $\{y^4, y^d\}$, $\{y^5, y^c\}$, $\{y^6, y^b\}$, $\{y^7, y^a\}$ and $\{y^8, y^9\}$, where $y \in \{A, B, C, D, E, F\}$. For example, $\aleph(A, D, C, F)$ and $\aleph(A^0, D^0, C^0, F^0)$ are represented in Figure 1 below, where A, A^0 (resp. D, D^0; C, C^0; D, D^0 have thick black (resp. green; blue; red) edges.



Figure 1: Assemblies $\aleph(A, D, C, F)$ and $\aleph(A^0, D^0, C^0, F^0)$ of $\mathcal{S}$

There is a copy of the dihedral group D_8 as a subgroup of the automorphism group $\mathcal{A}$ of $\mathcal{S}$ that has: **(a)** a cyclic subgroup of order 8 generated by an automorphism that sends A_i onto D_{2i} (given below by a matrix ρ) and **(b)** an automorphism that sends A_i onto A^0_i (given below by a matrix α^2), where $i \in \mathbf{Z}_{17}$. Moreover, $\mathcal{A}$ contains a subgroup $\mathcal{A}_0$ which is a semidirect product of this copy of D_8 and a copy of $\mathbf{Z}_{17}$ generated by a matrix θ, also given below.

There are 18 reconstructions of $\mathcal{S}$, of which we just treated $\aleph(A, D, C, F)$ and $\aleph(A^0, D^0, C^0, F^0)$. The remaining sixteen reconstructions, denoted $\aleph(A^j, D^j, C^j, F^j)$, have 17-cycles A^j, D^j, C^j, F^j, with $0 \neq j \in \mathbf{Z}_{17}$, obtained from A^0, D^0, C^0, F^0 via uniform addition of indexes mod 17. If we denote by $\mathcal{A}|\zeta$ the order-9 subgroup of $\mathcal{A}$ generated by an adequate rotation ζ of a 9-cycle of $\mathcal{S}$, then the 18 reconstructions above are related by means of the subgroups $\mathcal{A}|\zeta$. Each such subgroup $\mathcal{A}|\zeta$ of $\mathcal{A}$: **(a)** has a semidirect product with $\mathcal{A}_0$ equal to $\mathcal{A}$, showing that $|\mathcal{A}| = 17 \times 3^2 \times 2^4$ and how $\mathcal{A}$ is structured by means of semidirect products from D_8, $\mathbf{Z}_{17}$ and $\mathcal{A}|\zeta$; **(b)** partitions $V(\mathcal{S})$ into eleven 9-orbits and one 3-orbit $\mathcal{O}_\zeta$. The stabilizer in $\mathcal{A}|\zeta$ of any of the three vertices of $\mathcal{O}_\zeta \subset V(\mathcal{S})$ (on which $\mathcal{A}|\zeta \subset \mathcal{A}$ acts) contains a copy of the symmetric group S_3. Each one of these three vertices have distance 3 from ζ realized by paths whose ends in ζ subdivide it into paths of length 3. (This disposition of vertices will yield in Section 4 three disjoint copies of K_4 in $\mathcal{Y}$). For example, if $\zeta = (F_g F_8 F_0 F_9 F_1 E_1 D_1 D_g E_g)$, then $\mathcal{O}_\zeta = (A_0\, D_7\, D_a)$.



Figure 2: Frucht diagrams of $\mathcal{S}$ via $\mathbf{Z}_{17}$ and $\mathbf{Z}_9$

There is an 11-arc departing from each vertex of ζ, yielding a total of 9 disjoint 11-arcs, three arriving at each vertex of $\mathcal{O}_\zeta$, with component vertices at each fixed distance $d < 11$ from ζ forming a corresponding 9-orbit $\mathcal{O}^d$, so we can write $\mathcal{O}^0 = \zeta$ and $\mathcal{O}^{12} = \mathcal{O}_\zeta$. A corresponding Frucht diagram [13] of $\mathcal{S}$ via $\mathbf{Z}_9$ is depicted on the right of Figure 2, where each number denoting a vertex is followed with the parenthesized size of the $\mathbf{Z}_9$-orbit it represents, and with nonzero Frucht notation [13] over edges taken as arcs from left to right, so that the corresponding oppositely oriented arcs have the respective negative Frucht notation; horizontal edges, not having an explicit Frucht notation denoted, must be taken with null Frucht notation. There is a correspondence between the original notation of $\mathcal{S}$ and a new one based on this Frucht diagram, obtained by re-expressing the 9 11-arcs as follows:

$$\begin{array}{lcl}
F_g F_7 F_f E_f D_f D_d E_d B_d C_d C_0 B_0 A_0 & \rightarrow & 0_0\, 1_0\, 2_0\, 3_0\, 4_0\, 5_0\, 6_0\, 7_0\, 8_0\, 9_0\, a_0\, b_0 \\
F_8 E_8 B_8 A_8 A_9 A_a A_b B_b E_b D_b D_9 D_7 & \rightarrow & 0_1\, 1_1\, 2_1\, 3_1\, 4_1\, 5_1\, 6_1\, 7_1\, 8_1\, 9_1\, a_1\, b_1 \\
F_0 E_0 D_0 D_2 E_2 B_2 C_2 C_6 C_a B_a E_a D_a & \rightarrow & 0_2\, 1_2\, 2_2\, 3_2\, 4_2\, 5_2\, 6_2\, 7_2\, 8_2\, 9_2\, a_2\, b_2 \\
F_9 E_9 B_9 C_9 C_5 B_5 A_5 A_4 A_3 A_2 A_1 A_0 & \rightarrow & 0_3\, 1_3\, 2_3\, 3_3\, 4_3\, 5_3\, 6_3\, 7_3\, 8_3\, 9_3\, a_3\, b_0 \\
F_1 F_a F_2 F_b\; F_3 F_c\; F_4\; F_d F_5 E_5\; D_5 D_7 & \rightarrow & 0_4\, 1_4\, 2_4\, 3_4\, 4_4\, 5_4\, 6_4\, 7_4\, 8_4\, 9_4\, a_4\, b_1 \\
E_1 B_1 C_1 C_e B_e A_e A_d A_c B_c E_c D_c D_a & \rightarrow & 0_5\, 1_5\, 2_5\, 3_5\, 4_5\, 5_5\, 6_5\, 7_5\, 8_5\, 9_5\, a_5\, b_2 \\
D_1 D_3 E_3 B_3 C_3 C_7 C_b C_f B_f A_f A_g A_0 & \rightarrow & 0_6\, 1_6\, 2_6\, 3_6\, 4_6\, 5_6\, 6_6\, 7_6\, 8_6\, 9_6\, a_6\, b_0 \\
D_g D_e E_e F_e F_6 E_6 B_6 A_6 A_7 B_7 E_7 D_7 & \rightarrow & 0_7\, 1_7\, 2_7\, 3_7\, 4_7\, 5_7\, 6_7\, 7_7\, 8_7\, 9_7\, a_7\, b_1 \\
E_g B_g C_g C_c C_8 C_4 B_4 E_4 D_4 D_5 D_8 D_a & \rightarrow & 0_8\, 1_8\, 2_8\, 3_8\, 4_8\, 5_8\, 6_8\, 7_8\, 8_8\, 9_8\, a_8\, b_2
\end{array}$$

The 18 reconstructions of $\mathcal{S}$ cited above correspond in a 1-1 fashion to the 18 points of the projective line $\mathcal{P}$ over the field $GF(17)$; this 1-1 correspondence is to be set by the end of this section. Elements of $PSL(2,17)$, represented by 2×2-matrices over $GF(17)$, act over these points by left multiplication. In order to fit with the 2×2-matrices provided below, $\aleph(A,D,C,F)$ must correspond to the point of $\mathcal{P}$ given by the column vector $\binom{1}{9}$. Then the point at ∞, given by $\binom{0}{1}$, and the remaining points, given by vector columns $\binom{1}{j}$, for $9\neq j\in GF(17)$, correspond to the remaining reconstructions $\aleph(A^i,D^i,C^i,F^i)$, for $i\in GF(17)$.

The stabilizer of the group $PSL(2,17)$ at any fixed element of $\mathcal{P}$ coincides with the automorphism group of the 3-cube Q_3. To see that this is also the stabilizer $\mathcal{A}_{A_0}$ of $\mathcal{A}$ at the vertex A_0 of $\mathcal{S}$, recall that the number of 4-arcs departing from a fixed vertex of $\mathcal{S}$ is 24. Each such 4-arc determines a unique oriented 17-cycle and a unique oriented 9-cycle of $\mathcal{S}$. Thus, there are twenty four oriented 17-cycles passing through each vertex of $\mathcal{S}$, in particular through A_0. Two of these oriented 17-cycles are $A=(A_0A_1A_2\dots A_g)$ and its oppositely oriented 17-cycle, $-A$. The other oriented 17-cycles through A_0 are obtained from the reconstructions of $\mathcal{S}$ above. In fact, $\mathcal{A}_{A_0}$ has generators α,β,γ of orders 4, 2 and 3 respectively, that map the successive vertices of A, as presented above, onto the respective successive vertices of $-D^9, D^{10}, -C^3$. Moreover, α,β,γ form a generating set of the automorphism group of Q_3. For example, the automorphism that sends A onto $-A$ equals $\alpha^2\beta\gamma^2$. Likewise, A is sent onto the following 17-cycles, by means of the automorphisms expressed in the subsequent contiguous line:

$$\begin{array}{lllllllllll}
A^0 & A^1 & C^3 & C^4 & D^7 & D^8 & D^9 & D^a & C^d & C^e & A^g \\
\alpha^2 & \alpha^3\gamma & \alpha^3\gamma^2 & \alpha\gamma & \alpha\gamma^2 & \alpha^3 & \alpha^3\beta\gamma^2 & \beta & \alpha^2\beta\gamma & \alpha^2\beta & \beta\gamma \\
 & & & & & & & & & & \\
-A^0 & -A^1 & -C^3 & -C^4 & -D^7 & -D^8 & -D^9 & -D^a & -C^d & -C^e & -A^g \\
\beta\gamma^2 & \alpha^2\gamma^2 & \gamma & \gamma^2 & \alpha^2\gamma & \alpha\beta\gamma^2 & \alpha & \alpha\beta\gamma & \alpha\beta & \alpha^3\beta\gamma & \alpha^3\beta
\end{array}$$

completing the specifications for the 24 elements of $\mathcal{A}_{A_0}$, that yield information about 12 of the 18 reconstructions of $\mathcal{S}$, with each two oppositely oriented 17-cycles displayed belonging to a specific reconstruction. These twelve reconstructions are respectively $\aleph(A^i,D^i,C^i,F^i)$, for $i=0,1,3,4,7,8,9,a,d,e,g$.

The generators α,β,γ of $\mathcal{A}_{A_0}$ can be associated with the following matrices:

$$\alpha\to\begin{pmatrix}0&1\\-1&6\end{pmatrix},\qquad \beta\to\begin{pmatrix}0&1\\-1&0\end{pmatrix},\qquad \gamma\to\begin{pmatrix}0&1\\-1&-1\end{pmatrix}.$$

These matrices allow to establish a group embedding of $\mathcal{A}_{A_0}$ onto $PSL(2,17)$. By Lagrange's theorem, $\mathcal{A}_{A_0}$ partitions $\mathcal{A}$ into 102 classes, corresponding to the 102 vertices of $\mathcal{S}$. Also, γ has as cubic roots the following three matrices:

$$(\sqrt[3]{\gamma})_1\to\begin{pmatrix}2&8\\9&11\end{pmatrix},\qquad (\sqrt[3]{\gamma})_2\to\begin{pmatrix}6&15\\2&8\end{pmatrix},\qquad (\sqrt[3]{\gamma})_3\to\begin{pmatrix}9&11\\6&15\end{pmatrix}.$$

Any of these cubic roots can be associated to a rotation generator of $\mathcal{A}|\zeta$. Let $(\sqrt[3]{\gamma})_1$ be such a generator. We can take the copies of the subgroups $\mathbf{Z}_{17}$ and D_8 in $\mathcal{A}$ above to be generated in $PSL(2,17)$ by its elements represented by

$$\theta\to\begin{pmatrix}0&2\\8&2\end{pmatrix},\qquad \rho\to\begin{pmatrix}0&1\\-1&8\end{pmatrix},\qquad \alpha^2\to\begin{pmatrix}-1&6\\-6&1\end{pmatrix},$$

where θ has order 17, ρ has order 8 and ρ combines with α^2 (or with β) to form a copy of D_8 in $\mathcal{A}$. These copies of the two subgroups $\mathbf{Z}_{17}$ and D_8 in $\mathcal{A}$, related adequately via semidirect products to the copy of $\mathbf{Z}_9$ generated by $(\sqrt{\gamma})_1$, yields an isomorphism $\mathcal{A} \rightarrow PSL(2,17)$ to be used specifically in Theorem 7.

Because A and $-A$ are represented by the identity and by $\alpha^2\beta\gamma^2$, respectively, it can be seen that the reconstruction $\aleph(A, D, C, F)$ of $\mathcal{S}$ has associated the point $\binom{1}{9}$ of $\mathcal{P}$ with the vertex A_0. Similarly, because A^0 and $-A^0$ are represented by α^2 and by $\beta\gamma^2$, respectively, the reconstruction $\aleph(A^0, D^0, C^0, F^0)$ of $\mathcal{S}$ has associated the point $\binom{1}{10}$ of $\mathcal{P}$ with the vertex A_0. The selection of θ is motivated for it takes this point, $\binom{1}{10} \in \mathcal{P}$, via successive powers $\theta, \theta^2, \ldots$ onto the following points:

$$\tbinom{1}{10}, \tbinom{1}{15}, \tbinom{1}{16}, \tbinom{1}{14}, \tbinom{1}{11}, \tbinom{1}{6}, \tbinom{1}{13}, \tbinom{1}{0}, \tbinom{0}{1}, \tbinom{1}{1}, \tbinom{1}{5}, \tbinom{1}{12}, \tbinom{1}{7}, \tbinom{1}{4}, \tbinom{1}{2}, \tbinom{1}{3}, \tbinom{1}{8},$$

which correspond to the successive reconstructions $\aleph(A^i,\ D^i,\ C^i,\ F^i)$ of $\mathcal{S}$, for $i = 0, \ldots, 8, 10, \ldots, 16$, thus establishing the desired 1-1 correspondence from the family of 18 reconstructions of $\mathcal{S}$ onto $\mathcal{P}$.

Theorem 3 *In $\mathcal{S}$, there exist* 18 *disjoint unions of four* 17*-cycles each, determining* 18 *corresponding reconstructions of $\mathcal{S}^3$. Those unions in $\mathcal{S}$ are in* 1-1 *correspondence with the* 18 *points of the projective line $\mathcal{P}$ of $GF(17)$ in such a way that the action of $\mathcal{A}$ on the family of these reconstructions is equivalent to the natural action of $PSL(2,17)$ on $\mathcal{P}$.* □

4 The $\{K_4, L(Q_3)\}_{K_3}$-UH graph $\mathcal{Y} = \mathcal{S}^3$

We keep using the notation of $\mathcal{S}$ and its collection $\mathcal{C}_9$ of 136 9-cycles in Section 3. Let $(\mathcal{C}_9)^3$ be the collection of distance-3 graphs of 9-cycles of $\mathcal{C}_9$. In each copy w_0w_1 of P_2 in a member $C_9^3 = (C_9)^3$ of $(\mathcal{C}_9)^3$, where $C_9 \in \mathcal{C}_9$, let us denote by e the edge between w_0 and w_1. Then, the initial vertex w_0, the initial flag $\{w_0, e\}$, the terminal flag $\{e, w_1\}$ and the terminal vertex w_1 of w_0w_1 are indicated, respectively, by means of the names of the vertices v_0, v_1, v_2, v_3 of the copy $v_0v_1v_2v_3$ of P_4 in C_9 for which w_0w_1 stands in C_9^3.

For example, if $C_9 = S^0 = (A_0A_1B_1C_1C_5C_9C_dC_0B_0)$, so that $C_9^3 = (S^0)^3 = (A_0C_1C_d)(A_1C_5C_0)(B_1C_9B_0)$, then the initial flag of the copy A_0C_1 of P_2 in $C_9^3 = (S^0)^3$ is indicated by A_1, the terminal flag by B_1, while A_0 and C_1 are indicated by themselves, namely A_0 and C_1. We get the indications over $C_9^3 = (S^0)^3$ shown in Figure 3.

We *zip* now corresponding arc pairs of the distance-3 graphs C_9^3 obtained from $\mathcal{S}$, in order to get a graph $\mathcal{Y}$ with the desired $\mathcal{C}$-UH properties, meaning that



Figure 3: Example of indications over $C_9^3 = (S^0)^3 = (A_0A_1B_1C_1C_5C_9C_dC_0B_0)^3$

we identify into an edge corresponding oppositely oriented arcs of the resulting 3-cycles. The following sequence of operations is performed; (compare with [7]):

$$\mathcal{S} \;\to\; \mathcal{C}_9 \;\to\; (\mathcal{C}_9)^3 \;\to\; \mathcal{Y}. \tag{1}$$

Next, we explain how this operation $\mathcal{S} \to \mathcal{Y}$ is composed. The distance-3 graphs C_9^3 of the 136 9-cycles C_9 of $\mathcal{S}$ are formed by three disjoint 3-cycles each, yielding a total of $3 \times 136 = 408$ 3-cycles. Thus, $\mathcal{C}_9$ determines a collection of 408 triangles in $\mathcal{Y}$, with each edge of $\mathcal{Y}$ shared by exactly two such triangles. In fact, we identify or zip in these triangles the pairs of copies of P_2 obtained as distance-3 graphs of copies of P_4 in $\mathcal{S}$. This yields 102 copies of K_4 that can be subdivided into six subcollections $\{Y^i\}$ of 17 copies of K_4 each, where $Y \in \{A,B,C,D,E,F\}$ and $i \in \{0,1,\ldots,16=g\} = \mathbf{Z}_{17}$. The vertex sets $V(Y^i)$, each followed by the set $\Lambda(Y_i)$ of copies of K_4 containing the corresponding vertex Y_i (as in the notation of Section 3), can be taken as follows, showing $\mathbf{Z}_2$-symmetries produced by the change-of-sign involution in $\mathbf{Z}_{17}$:

$$\begin{array}{llllllll}
V(A^i)=\{C_i, & D_i, & E_{i+4}, & E_{i-4}\}, & \Lambda(A_i)=\{C^i, & D^i, & E^{i+7}, & E^{i-7}\},\\
V(B^i)=\{D_{i+3}, & D_{i-3}, & F_{i+5}, & F_{i-5}\}, & \Lambda(B_i)=\{D^{i+2}, & D^{i-2}, & F^{i+8}, & F^{i-8}\},\\
V(C^i)=\{A_i, & F_i, & E_{i+1}, & E_{i-1}\}, & \Lambda(C_i)=\{A^i, & F^i, & E^{i+6}, & E^{i-6}\},\\
V(D^i)=\{A_i, & D_i, & B_{i+2}, & B_{i-2}\}, & \Lambda(D_i)=\{A^i, & D^i, & B^{i+3}, & B^{i-3}\},\\
V(E^i)=\{C_{i+6}, & C_{i-6}, & A_{i+7}, & A_{i-7}\}, & \Lambda(E_i)=\{C^{i+1}, & C^{i-1}, & A^{i+4}, & A^{i-4}\},\\
V(F^i)=\{C_i, & F_i, & B_{i+8}, & B_{i-8}\}, & \Lambda(F_i)=\{C^i, & F^i, & B^{i+5}, & B^{i-5}\},
\end{array}$$

where $i \in \mathbf{Z}_{17}$. This reveals a duality map ϕ from the 102 vertices of $\mathcal{S}$ onto the 102 copies of K_4. In fact, the copies of K_4 are the vertices

$$\begin{array}{lll}
\phi(A_i) = A^{3i} = A^*_i, & \phi(B_i) = B^{-7i} = B^*_i, & \phi(C_i) = C^{3i} = C^*_i,\\
\phi(D_i) = D^{5i} = D^*_i, & \phi(E_i) = E^{6i} = E^*_i, & \phi(F_i) = F^{5i} = F^*_i,
\end{array}$$

($i \in \mathbf{Z}_{17}$), of a graph $\phi(\mathcal{S}) = \mathcal{S}^* \equiv \mathcal{S}$ with a structure similar to that of the vertices $A_i, B_i, C_i, D_i, E_i, F_i$ of $\mathcal{S}$ and whose copies of K_4 can be precisely denoted $y_i = A_i, B_i, C_i, D_i, E_i, F_i$, with corresponding vertex sets $\Lambda(y_i)$ as specified above. Moreover, $\phi : \mathcal{S} \to \mathcal{S}^*$ is a graph isomorphism, with the adjacency of $\mathcal{S}^*$ similar to that of $\mathcal{S}$.

The distance-3 graph of each 9-cycle of $\mathcal{S}$ is composed by three disjoint 3-cycles in $\mathcal{Y}$. For example, the distance-3 graphs of $S^0, \ldots, Z^0$ are:

$$\begin{array}{l}
S^0 \to \{E^7\backslash A_e=(A_0,C_1,C_d),\ E^b\backslash A_4=(A_1,C_5,C_0),\ F^9\backslash F_9=(B_1,C_9,B_0)\};\\
T^0 \to \{E^b\backslash C_5=(C_0,A_4,A_1),\ E^a\backslash C_g=(C_4,A_3,A_0),\ D^2\backslash D_2=(B_4,A_2,B_0)\};\\
U^0 \to \{C^1\backslash A_1=(E_0,F_1,E_2),\ B^5\backslash D_8=(F_0,F_a,D_2),\ B^e\backslash D_b=(F_9,F_2,D_0)\};\\
V^0 \to \{A^4\backslash C_4=(E_0,D_4,E_8),\ B^3\backslash F_f=(D_0,D_6,F_8),\ B^5\backslash F_a=(D_2,D_8,F_0)\};\\
W^0 \to \{C^0\backslash E_g=(A_0,E_1,F_0),\ C^1\backslash E_2=(A_1,F_1,E_0),\ F^9\backslash C_9=(B_1,F_9,B_0)\};\\
X^0 \to \{A^0\backslash E_d=(C_0,E_4,D_0),\ A^4\backslash E_8=(C_4,D_4,E_0),\ D^2\backslash A_2=(B_4,D_2,B_0)\};\\
Y^0 \to \{C^1\backslash F_1=(E_0,A_1,E_2),\ D^2\backslash B_4=(B_0,A_2,D_2),\ D^0\backslash B_f=(A_0,B_2,D_0)\};\\
Z^0 \to \{F^0\backslash B_9=(F_0,B_8,C_0),\ F^8\backslash B_g=(F_8,C_8,B_0),\ A^4\backslash D_4=(E_8,C_4,E_0)\}.
\end{array}$$

This way, it can be seen that $\mathcal{Y}$ is a K_4-UH graph. Moreover, the vertices and copies of K_4 in $\mathcal{Y}$ are the points and lines of a self-dual (102_4)-configuration, which in turn has $\mathcal{Y}$ as its Menger graph. However, in view of Beineke's characterization of line graphs [1], and observing that $\mathcal{Y}$ contains induced copies of $K_{1,3}$, which are forbidden for line graphs of simple graphs, we conclude that $\mathcal{Y}$ is non-line-graphical. We obtain the following statement.

Theorem 4 *$\mathcal{Y}$ is an edge-disjoint union of 102 copies of K_4, with four such copies incident to each vertex. Moreover, $\mathcal{Y}$ is a non-line-graphical K_4-UH graph. Its vertices and copies of K_4 are the points and lines, respectively, of a self-dual (102_4)-configuration which has precisely $\mathcal{Y}$ as its Menger graph. In particular, $\mathcal{Y}$ is arc-transitive with regular degree 12, diameter 3 and distance distribution $(1, 12, 78, 11)$. Its associated Levi graph* [5] *is a 2-arc-transitive graph with regular degree 4, diameter 6 and distance distribution $(1, 4, 12, 36, 78, 62, 11)$.*

Proof. The statement is immediate from the data given by the construction of $\mathcal{Y}$ via operation (1) and because of the fact that $\mathcal{S}$ is a distance-transitive graph. $\square$

Remark 5 *A proof yielding the group structure of the automorphism group $\mathcal{B}$ of $\mathcal{Y}$ is part of the proof of Theorem 6. However, the Magma software allowed to verify that the automorphism group of $\mathcal{Y}$ is $\mathcal{B} = \mathcal{A} = PLS(2, 17)$, whose order is 2448, and that the Levi graph associated to $\mathcal{Y}$ has automorphism group $SL(2, 17)$, whose order is 4896.*

Theorem 6 *$\mathcal{Y}$ is a $\{K_4\}$-UH graph.*

Proof. Consider an isomorphism $\Psi : \Theta_1 \to \Theta_2$ between two copies Θ_1, Θ_2 of K_4 in $\mathcal{Y}$. By construction, each Θ_i $(i = 1, 2)$ arises from 4 9-cycles θ_i^j in $\mathcal{S}$ $(j = 1, 2, 3, 4)$ related so that each one of them have their union as a subgraph $\overline{\Theta}_i$ of $\mathcal{S}$ with 4 vertices v_i^j of degree 3 and 12 vertices of degree 2 which are the internal vertices of 6 3-paths whose endvertices are the vertices v_i^j. For example, the vertices $v_1^1 = B_0, v_1^2 = B_1, v_1^3 = F_9, v_1^4 = C_9, v_2^1 = B_1, v_2^2 = B_2, v_2 1^3 = F_a, v_2^4 = C_a$ in $\mathcal{S}$ determine such subgraphs Θ_1, Θ_2 in $\mathcal{Y}$ and $\overline{\Theta}_1, \overline{\Theta}_2$ in $\mathcal{S}$. Clearly, Ψ induces an isomorphism $\overline{\Psi} : \overline{\Theta}_1 \to \overline{\Theta}_2$ that sends say each v_1^j onto a corresponding v_2^j $(j = 1, 2, 3, 4)$. An automorphism $\overline{\overline{\Psi}}$ of $\mathcal{S}$ exists that extends $\overline{\Psi}$. Moreover, $\overline{\overline{\Psi}}$ determines an automorphism of $\mathcal{Y}$ that restricts to Ψ, which completes our verification that $\mathcal{Y}$ is a $\{K_4\}$-UH graph. $\square$

Each of the 102 copies of K_4 in $\mathcal{Y}$ arises from the distance-3 graphs of four of the 136 9-cycles of $\mathcal{S}$. The subgraph of $\mathcal{S}$ spanned by these four 9-cycles contains four degree-3 vertices, (which are initial and terminal vertices of corresponding copies of P_4, as in the first paragraph of the section), and twelve degree-2 vertices, (internal vertices in those copies of P_4). These twelve vertices form a copy $\mathcal{L}$ of $L(Q_3)$ in $\mathcal{Y}$. For the copy A^0 of K_4 in $\mathcal{Y}$, the copy $\mathcal{L} = a^0$ of $L(Q_3)$ in $\mathcal{Y}$ can



Figure 4: Copy a^0 of $L(Q_3)$ in $\mathcal{Y}$

be represented as in the long rectangle $\mathcal{R}$ in Figure 4, where: **(a)** the leftmost and rightmost dotted lines of $\mathcal{R}$ must be identified by parallel translation; **(b)** each of the 8 shown copies Δ of K_3 forms part of a corresponding copy ∇ of K_4 (among the 102 in $\mathcal{Y}$) with its name written on the exterior of $\mathcal{R}$ about the horizontal edge of Δ and with the fourth vertex of ∇ having its name written on the interior of Δ; **(c)** the edges are colored via an edge partition Ω into 2-paths, each having its two composing edges displaying a common color from a set of three colors that are present in every triangle, with equally colored opposite edges in every 4-hole, where red, green and blue edges are indicated respectively with color symbols 1, 2 and 3. For $y = a, b, c, d, e, f$, we denote the copies y^0 of $L(Q_3)$ as follows:

a^0: $(D_fD_4C_4B_d)$ $(B_4C_dD_dD_2)$ $F_dE_0F_4B_0$ $(B^1D_4A^4C_4F^4B_dD^fD_f)$ $(F^dB_5A^dE_9B^gF_bD^2A_2)$;
b^0: $(D_5D_gE_cF_d)$ $(D_cF_4E_5D_1)$ $F_eE_3E_eF_3$ $(B^2F_7A^gC_gC^dA_dB^8D_b)$ $(B^9D_6C^4A_4A^1C_1B^aD_f)$;
c^0: $(F_8F_1A_1B_g)$ $(B_1A_gF_gF_9)$ $D_gE_0D_1B_0$ $(B^dD_aC^1E_2D^1B_3F^8C_8)$ $(D^gB_eC^gE_fB^4D_7F^9C_9)$;
d^0: $(A_1A_fD_fE_2)$ $(E_fD_2A_2A_g)$ $C_2B_0C_fE_0$ $(E^8C_eD^fB_dA^fE_bC^1F_1)$ $(A^2E_6D^2B_4E^9C_3C^gF_g)$;
e^0: $(A_6A_9B_bC_2)$ $(A_bC_fB_6A_8)$ $C_aB_7B_aC_7$ $(E^gC_5D^9D_9F^2F_2E^dA_3)$ $(E^4A_eF^fF_fD^aE_8E^1C_c)$;
f^0: $(C_4C_9F_9E_8)$ $(E_9F_8C_8C_d)$ $A_8B_0A_9E_0$ $(E^fA_5F^9B_1C^9E_aA^4D_4)$ $(C^8E_7F^8B_gE^2A_cA^dD_d)$;

and denote y^i via translation mod 17, for $0 \neq i \in \mathbf{Z}_{17}$. These a^0, c^0, d^0, f^0, (resp. b^0, e^0) are similar, say with respect to D_i, F_i, A_i, C_i, (resp. E_i, B_i), etc. Each copy y^i of $L(Q_3)$ admits an edge partition $\Omega = \Omega(y^i)$ into j-colored 2-paths ($j \in \{1, 2, 3\}$ as in Figure 4) in such a way that each (monochromatic) 2-path in an $\Omega(y^i)$ is shared only by another copy of $L(Q_3)$ in $\mathcal{Y}$, (see item **(3)** of Theorem 7, below). We may write $y^i = y^i_1 \cup y^i_2 \cup y^i_3$, the vertex-spanning edge-disjoint union of the red, green and blue subgraphs in each case, as is the situation in Figure 4 for $y^i = a^0$.

Each vertex of $\mathcal{Y}$ belongs exactly to twelve such copies $\mathcal{L}$ of $L(Q_3)$. Figure 5 shows the complements of A_0 in four of the twelve copies of $L(Q_3)$ containing A_0 (sharing the long vertical edges), where the black vertices form the 8 4-holes containing A_0, and some edges and vertices appear twice, in fact once per copy of $L(Q_3)$. For example, the leftmost and rightmost edges must be identified by parallel translation. Alternate internal anti-diagonal (i.e. from top-right to bottom-left) 2-paths in Figure 5 also coincide with their directions reversed. This contributes to the edge partitions $\Omega(y^i)$ into 2-paths, exactly the



Figure 5: Complements of A_0 in four of the twelve copies of $L(Q_3)$

2-paths along which these coincidences occur. (Notice that the middle vertices of these 2-paths are the neighbors of A_0 in $\mathcal{S}$, and that their degree-1 vertices are at distance 2 from A_0 in $\mathcal{S}$). The 9-cycles of the $\mathcal{C}_9$ in Section 3 intervene in the formation of the oriented 3-cycles containing A_0, (namely $E^7 \setminus A_e = (A_0, C_1, C_d) \subset S^0$, $E^a \setminus C_g = (A_0, B_2, D_0) \subset Y^0$, $D^0 \setminus B_f = (A_0, B_2, D_f) \subset Y^0$, $C^0 \setminus E_g = (A_0, E_1, F_0) \subset W^0$, all cited in the table displayed shortly before Theorem 3, and of which Figure 5 just mentions the corresponding containing copies of K_4: E^7, E^a, D^0, C^0) and the involved copies of $L(Q_3)$ (namely e^b, d^2, c^1, f^9), induced respectively by the long vertical edges (namely C_dC_1, A_3C_4, D_0B_2, E_1F_0) and by the 6-cycles they separate (namely $(C_dB_5A_3C_4A_5C_1)$, $(A_3E_4D_0B_2D_4C_4)$, $(D_0F_2E_1F_0F_aB_2)$, $(E_1E_9C_dC_1B_9F_0)$). There are two other complements of A_0 in the union of four copies of $L(Q_3)$, yielding a total of three, whose symbolic information can be set as in the following three respective arrays, the left one summarizing the data in Figure 5, where the two twice-repeated middle vertices appear between parentheses to the right of the heading A_0 and the remaining data displayed as in Figure 5:

$$\begin{array}{ccc}
A_0(B_0A_1) & A_0(A_1A_g) & A_0(A_gB_0) \\
(E^7e^bE^ad^2D^0c^1C^0f^9) & (E^7d^gC^0d^1E^ae^9D^0e^8) & (E^7d^fD^0c^gC^0f^8E^ae^6) \\
(C_dB_5A_3E_4D_0F_2E_1E_9) & (C_1E_eE_gD_3A_3C_7B_fC_6) & (A_eE_dD_0F_fE_gE_8C_4B_c) \\
(B_4B_1E_2C_0F_9A_2C_5E_0) & (D_1A_fC_3B_1C_2B_gB_eA_2) & (E_fC_0F_8A_fC_cE_0B_dB_g) \\
(A_2C_9E_0A_4B_1D_2C_0F_1) & (B_gC_eA_2D_gA_fB_3B_1C_f) & (E_0A_dB_gD_fC_0F_gA_fC_8) \\
(C_1A_5C_4D_4B_2F_aF_0B_9) & (A_eD_eE_1E_3C_gC_bB_2C_a) & (C_dD_dB_fF_7F_0B_8C_gA_c)
\end{array}$$

Some edges are shared by two different of these three arrays. In fact, each of the edges bordering the 2-paths ω presented in anti-diagonal 4-paths as in Figure 5 is present also in the second or third arrangement. For example, the edge B_1A_3 of E^4 on Figure 5 appears in the second arrangement. The vertices B_0, C_0, D_0, E_0 and F_0 admit similar arrangements. Additions mod 17 yield the remaining information for neighboring copies of K_4 and $L(Q_3)$ at each vertex of $\mathcal{Y}$. From the facts presented up to this point, we have the following properties expressed as a single theorem.

Theorem 7 **(1)** *$\mathcal{Y}$ is a connected union of* 102 *copies of $L(Q_3)$, each with an edge partition Ω into 2-paths.* **(2)** *Each edge in $\mathcal{Y}$ is shared exactly by four copies of $L(Q_3)$ in $\mathcal{Y}$.* **(3)** *Each copy Δ of K_3 (resp. 2-path $\omega \in \Omega(y)$) of a copy y of $L(Q_3)$ in $\mathcal{Y}$ is shared exactly by two copies y, y' of $L(Q_3)$ in $\mathcal{Y}$.* **(4)** *Each two copies of $L(Q_3)$ sharing a copy Δ of K_3 in $\mathcal{Y}$ share Δ with exactly one copy* *of* K_4 *in* $\mathcal{Y}$. **(5)** *each* 4*-hole of* $\mathcal{Y}$ *exists in just one copy of* $L(Q_3)$ *in* $\mathcal{Y}$. **(6)** $\mathcal{Y}$ *is an* Ω*-preserving* SZ $\{L(Q_3)\}_{K_3}$-UH *graph.* **(7)** $\mathcal{Y}$ *is* $\{K_4, L(Q_3)\}_{K_3}$-UH. **(8)** *The automorphism group* $\mathcal{B}$ *of* $\mathcal{Y}$ *is* $\mathcal{A} = PSL(2,17)$.

In item **(3)** of Theorem 7, for each triangle Δ in y, the copies y, y' of $L(Q_3)$ intersect exactly in Δ, while for a 2-path $\omega \in \Omega(y)$ in y, not only ω is shared by y, y', but y, y' also share a vertex at distance 2 from the endvertices of ω. This common distance, 2, is realized by 2-paths in the other two colors $\neq$ from the color of ω, in each of y and y', as in Figure 4, where for example the red-colored 2-path $F_4D_2B_4$ (present both in a^0 and c^3) is at distance 2 from vertex D_4 (also present in a^0 and c^3) via the green-colored path $B_4F_dD_4$ and the blue-colored path $F_4C_4D_4$.

Proof. The $\mathcal{C}$-UH properties of $\mathcal{Y}$ in the statement arise because of its construction via operation (1) and the fact that $\mathcal{S}$ is distance-transitive.

We explain now how a 2-path-preserving isomorphism $\Psi' : \Theta'_1 \to \Theta'_2$ between two copies Θ'_1, Θ'_2 of $L(Q_3)$ in $\mathcal{Y}$ can be extended to an automorphism of $\mathcal{S}$. Both Θ'_1 and Θ'_2 are colored in the fashion of Figure 4 with Ψ' respecting colors, thus inducing a 1-1 correspondence between the color classes of Θ'_1 and Θ'_2. In each copy of $L(Q_3)$ in $\mathcal{Y}$ there are exactly 12 monochromatic 2-paths, 4 in each one of the 3 colors, and exactly 12 dichromatic 2-paths, a total of 24 2-paths. A 2-path-preserving isomorphism $\Psi' : \Theta'_1 \to \Theta'_2$ can be extended to an automorphism of $\mathcal{Y}$ because the information gathered in Θ'_i arises from corresponding information in a subgraph $\overline{\Theta'}_i$ in $\mathcal{S}$ ($i = 1,2$) so that Ψ' comes from an isomorphism $\overline{\Psi'} : \overline{\Theta'}_1 \to \overline{\Theta'}_2$. However, we notice that $\overline{\Theta'}_i = \overline{\Theta}_i$ ($i = 1,2$) for a corresponding Θ_i as in the proof of Theorem 4, but while the vertices of Θ'_i are the degree-2 vertices of $\overline{\Theta'}_i = \overline{\Theta}_i$, the vertices of Θ_i are the degree-3 vertices of $\overline{\Theta}_i = \overline{\Theta'}_i$. Here the pairs (Θ_i, Θ'_i) are of the form (Y^j, y^j), where $(Y, y) \in \{(A,a),(B,b),(C,c),(D,d),(E,e),(F,f)\}$ and $j \in \mathbf{Z}_{17}$. Then $\overline{\Psi'} = \overline{\Psi} : \Theta_1 \to \Theta_2$ is a corresponding map as in the proof of Theorem 4. But now $\overline{\Psi'} = \overline{\Psi}$ extends to an automorphism of $\mathcal{S}$, as in the proof of Theorem 4. This takes us to an automorphism of $\mathcal{Y}$ that extends Ψ', as claimed above.

For example, the red 2-path $B_4F_dD_4$ in the copy a^0 of $L(Q_3)$ in $\mathcal{Y}$ in Figure 4 is realized by the 3-paths $B_4E_4F_4F_d$ and $F_dF_4E_4D_4$ in $\mathcal{S}$, which share two edges with a common vertex but differ in an edge, with union of these 2 3-paths realized by a tree T_1 with just one vertex of degree 3, namely $t_1 = E_4$, from which 2 1-paths and one 2-path depart. A similar tree T_2 is obtained from the red 2-path $D_dF_4B_d$. This T_2 intersects T_1 on the 1-path F_dF_4, which is a terminal 2-path of both T_1 and T_2 on their 2-paths departing from $t_1 = E_4$ and degree 3-vertex $t_2 = E_d$ of T_2, respectively. The other two red 2-paths in Figure 4 behave similarly, leading to trees T_3 and T_4 intersecting at the 1-path B_0E_0. Similar behavior holds for the blue and the green 2-path quadruples in Figure 4, leading to pairs of trees that intersect respectively at the 1-paths D_4D_2, B_dC_d and the 1-paths B_4C_4, D_fD_d. Thus, if Θ'_1 is this copy of $L(Q_3)$ in $\mathcal{Y}$, then $\overline{\Theta'}_1$

coincides with $\overline{\Theta}_1$, where $\Theta_1 = A^0$. With this, the reader is invited to complete an example for the situation presented in the previous paragraph.

Now, we need only to prove item **(8)** of the statement of the theorem. Since we took the copies of P_2 in $\mathcal{Y}$ in correspondence with the copies of P_4 in $\mathcal{S}$, reattaching then the 3-cycles of $\mathcal{Y}$ resulting from the 9-cycles of $\mathcal{S}$ by zipping corresponding copies of K_2 in $\mathcal{Y}$, so leading to the 102 copies of K_4 and to the 102 copies of $L(Q_3)$), we have that: **(a)** the 17-cycles of $\mathcal{S}$ have their distance-3 graphs in $\mathcal{Y}$ as 17-cycles themselves, (for example, $(A_0A_1A_2\ldots A_g)$ in $\mathcal{S}$ is transformed into $(A_0A_3A_6\ldots A_e)$ in $\mathcal{Y}$, which uses the same vertices with a uniform adjacency advancement of three index units in each case, instead of one unit); **(b)** for each reconstruction $\aleph(P^1, P^2, P^4, P^8)$ of $\mathcal{S}$ as in Section 3, there is a reconstruction $\mathcal{A}'((P^1)^3, (P^2)^3, (P^4)^3, (P^8)^3)$ of $\mathcal{Y}$ determined by the distance-3 graphs $(P^1)^3$, $(P^2)^3$, $(P^4)^3$, $(P^8)^3)$ of P^1, P^2, P^4, P^8, respectively; **(c)** each $\mathcal{A}'((P^1)^3, (P^2)^3, (P^4)^3, (P^8)^3)$ has 8 automorphisms that form a cyclic subgroup of $\mathcal{B}$, obtained by multiplying the vertex indexes by 2 mod 17; **(d)** the index-translation action mod 17 on $\mathcal{S}$ appears multiplied by 3 mod 17 in $\mathcal{Y}$; in other words, an index-translation action holds on $\mathcal{Y}$ from a subgroup $\mathbf{Z}_{17}$ of $\mathcal{B}$. It can be seen that the structure of $\mathcal{A}$ presented in Theorem 2 is then preserved in $\mathcal{B}$. Since $|\mathcal{B}| = |\mathcal{A}|$, then $\mathcal{B}$ is isomorphic to $\mathcal{A}$. $\square$

The 2-paths ω mentioned before Theorem 7 form a partition of $\mathcal{Y}$ into 153 4-holes (not faces of $L(Q_3)$ in $\mathcal{Y}$); these can be obtained by adding $x \in \mathbf{Z}_{17}$ uniformly mod 17 to all indexes in the following generating-set table of such 4-holes, shown in the left column; the rightmost four pairs of copies of y^i_j (as in the item **(c)** that accompanies Figure 4) are shared by the subsequent 2-paths in each 4-cycle, and the vertex pair following the 4-cycle is formed by two vertices that alternately are at distance two from the endvertices of each composing 2-path:

$(A_2B_0B_1A_g)\ A_0A_1$	$(c^1_3\ e^b_2)$	$(e^7_2\ c^0_2)$	$(d^1_3\ e^8_3)$	$(e^a_3\ d^0_2)$
$(C_0A_gE_0A_1)\ A_0B_0$	$(d^f_2\ f^8_1)$	$(c^0_1\ d^0_1)$	$(d^2_3\ f^9_1)$	$(e^7_1\ e^a_1)$
$(C_4E_0C_dA_0)\ B_0C_0$	$(a^0_1\ f^0_1)$	$(f^9_2\ d^f_1)$	$(e^6_2\ e^b_2)$	$(d^2_1\ f^8_3)$
$(D_0A_0F_0C_0)\ B_0E_0$	$(c^g_2\ c^1_3)$	$(f^8_2\ f^9_3)$	$(a^d_2\ a^4_3)$	$(d^f_3\ d^2_2)$
$(C_8B_0B_4C_d)\ C_0C_4$	$(a^4_3\ e^a_1)$	$(e^b_1\ a^0_2)$	$(f^4_3\ e^f_3)$	$(e^6_3\ f^0_2)$
$(D_4D_fE_2E_0)\ D_0D_2$	$(a^2_2\ b^g_3)$	$(b^e_2\ d^0_3)$	$(d^2_2\ b^5_2)$	$(b^3_3\ a^0_3)$
$(F_0D_2B_0D_f)\ D_0E_0$	$(c^1_1\ a^4_2)$	$(a^0_3\ d^0_1)$	$(a^d_3\ c^g_1)$	$(b^3_1\ b^e_1)$
$(F_8B_0F_9D_0)\ E_0F_0$	$(c^0_1\ f^0_1)$	$(c^1_2\ a^d_1)$	$(b^5_2\ b^c_2)$	$(a^4_1\ c^g_3)$
$(E_8\ E_0F_gF_9)\ F_0F_8$	$(b^3_1\ f^8_2)$	$(b^c_3\ c^0_2)$	$(c^8_3\ b^d_3)$	$(f^0_3\ b^5_1)$

Since any vertex of each y^j_i has degree four in y^j_i, the corresponding $\Omega(y^j_i)$ has a well-defined complementary partition $\Omega'(y^j_i)$ of y^j_i into 3-paths. The collection of 3-paths in all the partitions $\Omega'(y^j_i)$ reassembles as a collection of 306 copies of $K_{1,4}$. A generating set of these copies is shown subsequently, with the remaining copies of $K_{1,4}$ obtained by addition of $x \in \mathbf{Z}_{17}$ uniformly mod 17 to all indexes $i \in \mathbf{Z}_{17}$ of vertices Y_i and subgraphs y^i_j, where $j = 1, 2, 3$ stand still for red, green and blue, respectively:

$A_0\,(e_3^b\,A_3\,d_2^1)\,(d_1^1\,E_1\,c_1^1)\,(c_2^1\,B_2\,e_3^8)\,(e_1^8\,C_1\,e_1^b)$	$D_0\,(b_1^c\,F_f\,b_1^1)\,(b_2^1\,E_d\,d_2^f)\,(d_1^f\,B_f\,a_1^f)\,(a_2^f\,D_b\,b_3^c)$
$A_0\,(f_3^8\,C_4\,d_1^2)\,(d_2^2\,D_0\,d_3^f)(d_1^f\,C_d\,f_2^9)\,(f_3^9\,F_0 f_2^8)$	$D_0\,(a_1^d\,F_9\,c_2^1)\,(c_3^1\,A_0\,c_2^g)\,(c_3^g\,F_8\,a_1^4)\,(a_3^4\,C_0\,a_2^d)$
$A_0\,(d_3^g\,A_e\,e_3^6)\,(e_1^6\,C_g\,e_1^9)\,(e_2^9\,B_f\,c_3^g)\,(c_2^g\,E_g\,d_1^g)$	$D_0\,(b_3^5\,D_6\,a_3^2)\,(a_1^2\,B_2\,d_1^2)\,(d_3^2\,E_4\,b_2^g)\,(b_1^g\,F_2\,b_1^5)$
$B_0\,(e_1^6\,B_d\,a_3^0)\,(a_2^0\,B_4\,e_1^b)\,(e_3^b\,C_9\,f_3^0)\,(f_2^0\,C_8\,e_3^6)$	$E_0\,(a_2^0\,D_d\,b_2^e)\,(b_2^e\,E_2\,d_3^0)\,(d_2^0\,E_f\,b_2^3)\,(b_3^3\,D_4\,a_3^0)$
$B_0\,(e_3^7\,A_f\,d_3^0)\,(d_2^0\,A_2\,e_3^a)\,(e_2^a\,B_g\,c_3^0)\,(c_2^0\,B_1\,e_2^7)$	$E_0\,(b_3^5\,F_1\,c_3^0)\,(c_2^0\,F_g\,b_3^c)\,(b_1^c\,E_9\,f_2^0)\,(f_3^0\,E_8\,b_1^5)$
$B_0\,(a_2^4\,D_2\,c_1^1)\,(c_2^1\,F_9\,a_1^d)\,(a_3^d\,D_f\,c_1^g)\,(c_3^g\,F_8\,a_1^4)$	$E_0\,(f_1^9\,A_1 d_3^2)\,(d_1^2 C_4\,f_3^8)\,(f_1^8\,A_g d_2^f)\,(d_1^f\,C_d\,f_2^9)$
$C_0\,(d_3^f\,D_0\,d_2^2)\,(d_3^2\,A_1 f_1^9)\,(f_3^9\,F_0\,f_2^8)\,(f_1^8 A_g d_2^f)$	$F_0\,(c_2^g\,A_0\,c_3^1)\,(c_1^1\,D_2\,a_2^4)\,(a_3^4\,C_0\,a_2^d)\,(a_3^d\,D_f\,c_1^g)$
$C_0\,(e_2^7\,A_d\,e_2^2)\,(e_1^2\,B_9\,a_3^d)\,(a_1^d\,E_d\,f_1^d)\,(f_3^d\,C_5 e_3^7)$	$F_0\,(b_2^d\,D_8\,b_2^3)\,(b_3^3\,F_7\,c_2^8)\,(c_1^8\,B_8\,f_1^8)\,(f_3^8\,E_g\,b_1^d)$
$C_0\,(d_1^f\,B_8\,a_2^4)\,(a_1^4\,E_4\,f_1^4)\,(f_2^4\,C_c\,e_3^a)\,(e_2^a\,A_4 d_2^f)$	$F_0\,(f_2^9\,E_1\,b_1^4)\,(b_2^4\,D_9\,b_2^e)\,(b_3^e\,F_a\,c_3^9)\,(c_1^9\,B_9\,f_1^9)$

This generating set of copies of $K_{1,4}$ has each entry starting with a degree-4 vertex Y_0 followed by four parenthesized expressions each containing as its central entry a neighbor Y' of Y_0 flanked by the two subgraphs y_j^i to which the edge Y_0Y' belongs, so that each participating y^i appears repeated twice (with two different colors j, j', as y_j^i and $y_{j'}^i$), once before a right parenthesis and once after the subsequent left parenthesis, where the first left parenthesis is considered as subsequent to the last right parenthesis.

Let $\mathcal{I}$ and $\mathcal{J}$ be the collections formed respectively by the 153 non-cuboctahedral 4-holes of $\mathcal{Y}$ and the just presented 306 copies of $K_{1,4}$. Then, $\mathcal{I}$ is an edge-disjoint collection of 4-holes in $\mathcal{Y}$ and $\mathcal{J}$ is a P_3-disjoint collection of copies of $K_{1,4}$ in $\mathcal{Y}$. Moreover, $\mathcal{Y}$ is $\mathcal{I}$-UH. However, we can only say that any homomorphism between members of $\mathcal{J}$ preserving the order of presentation of central-vertex neighbors in corresponding copies of $K_{1,4}$, as in the table above, extends to an automorphism of $\mathcal{Y}$.

Observe that each copy y of $L(Q_3)$ in $\mathcal{Y}$ intersects 8 other copies of $L(Q_3)$ in a triangle each, and twelve other copies of $L(Q_3)$ in a 2-path of $\Omega(y)$ plus one vertex at distance two from its endvertices each, a total of twenty copies of $L(Q_3)$ touching y either in a triangle Δ or in a 2-path ω of $\Omega(y)$ and the only isolated vertex at distance two from the endvertices of ω via monochromatic paths in the other two colors from that of ω.

The graph $\mathcal{I}'$ generated by the (diagonal) chords of the 4-cycles of $\mathcal{I}$ coincides with the distance-2 graph $\mathcal{S}^2$ of $\mathcal{S}$ and is an arc-transitive 6-regular graph of diameter 4 and automorphism group $PSL(2,17)$.

Expressing the copies of $K_{1,4}$ in $\mathcal{J}$ simply as $u(v)(w)(x)(y)$, (for example the copy of K_4 in the first line of the last table expressed simply as $A_0(A_3)(E_1)(B_2)(C_1)$), we may consider the graph $\mathcal{J}'$ generated by the corresponding 4-cycles (v, w, x, y). Then $\mathcal{J}'$ coincides with the distance-4 graph of $\mathcal{S}$ and is an arc-transitive 24-regular graph of diameter 2 and automorphism group $PSL(2, 17)$.

Now that we know that $\mathcal{I}' = \mathcal{S}^2$ and $\mathcal{J}' = \mathcal{S}^4$, we recognize finally that $\mathcal{Y} = \mathcal{S}^3$, (even though a similar result does not follow in the case of [7], since the Klein graph is not the distance-2 graph of the Coxeter graph).

Theorem 8 *$\mathcal{Y} = \mathcal{S}^3$, $\mathcal{I}' = \mathcal{S}^2$ and $\mathcal{J}' = \mathcal{S}^4$.*

5 Considerations related to Question 1

In trying to answer Question 1 in the Introduction, one may perform the Cartesian product of the graph $\mathcal{X}$ of [6] with a copy of K_2 to obtain a K_4-UH graph on 84 vertices, but this is not an edge-disjoint union of copies of K_4. Similarly, the rooted product of a K_4-UH vertex-transitive graph Λ with a K_4 is an edge-disjoint union of copies of K_4 and it is not a line graph if Λ is not a line graph, but it fails to be K_4-UH. More interesting are the Cartesian powers of K_4, $\mathcal{X}$ and $\mathcal{Y}$. These Cartesian powers are K_4-UH graphs but not Menger graphs of self-dual configurations. For example, the Cartesian square of K_4 is a K_4-UH graph as well as the Menger graph of a $(16_2, 8_4)$-configuration constituted by the lines $\{00, 01, 02, 03\}$, $\{10, 11, 12, 13\}$, $\{20, 21, 22, 23\}$, $\{30, 31, 32, 33\}$, $\{00, 10, 20, 30\}$, $\{01, 11, 21, 31\}$, $\{02, 12, 22, 32\}$ and $\{03, 13, 23, 33\}$, which is not a self-dual configuration. This situation is treated subsequently in more generality.

Theorem 9 *Let G be a connected K_4-UH graph that is not a line graph but is the edge-disjoint union of copies of K_4. Let H be the Cartesian product of G with itself, or Cartesian square of G. Then H is:* **(a)** *connected;* **(b)** *not a line graph;* **(c)** *the edge-disjoint union of copies of K_4;* **(d)** *K_4-UH.*

Proof. Item **(a)** arises because the Cartesian product of connected graphs is connected. Item **(b)** holds since H contains G as an induced subgraph but the class of line graphs is hereditary. Item **(c)** holds since in fact being a Cartesian product, H decomposes into edge-disjoint copies of G. For item **(d)**, note first that G is vertex transitive, since every vertex is incident with an edge, every edge is in a K_4 and since G is K_4-UH. By definition of the Cartesian product, for every K_4 in H one of the two natural projections to G maps the K_4 to only one vertex. Let A and B be two labeled copies of K_4 in H. We show that there is an automorphism that maps A to B. First note that the automorphism group of H contains the automorphism that interchanges the first factor with the second factor. Thus we may assume that both A and B project to a single vertex under the natural projection to the first factor. Suppose A maps to v_a and B to v_b under this projection. Since G is transitive there is an automorphism of G mapping v_a to v_b. This automorphism extends to an automorphism of H in the natural way. Thus we may assume that $v_a = v_b$. Let K_4^a and K_4^b be the (labeled) K_4 images of the natural projection to the second factor. There is an automorphism of G that maps K_4^a to K_4^b. This extends to an automorphism of H and maps A to B. □

Question 10 *Are there any non-line-graphical connected K_4-UH graphs that are edge-disjoint unions of copies of K_4, for positive integers n different from either* 4 *or* 42 *or* 102 *or a power of any of these three integers?*

Finally, [5] presents three self dual (n_4)-configurations, namely (5_4), (8_4) and and (4_4), whose Menger graphs are K_4-UH but not edge-disjoint unions of copies of K_4. They are obtained respectively from their Levi graphs, namely K_5, Q_4 and the result of adding to Q_4 the four edges between opposite vertices.

Acknowledgement. I thank the two referees of a previous version of this paper for their unselfish help with indications for correction and completion of proofs.

References

[1] L. W. Beineke, *Derived graphs and digraphs*, in Beiträge zum Graphentheorie, Teubner (1968) 17–33

[2] N. L. Biggs and D. H. Smith, *On trivalent graphs*, Bull. London Math. Soc., **3** (1971), 155-158.

[3] P. J. Cameron, 6-*transitive graphs*, J. Combin. Theory Ser. B **28** (1980), 168-179.

[4] I. Z. Bouwer et al., The Foster Census, R. M. Foster's Census of Connected Symmetric Trivalent Graphs, Charles Babbage Res. Ctr., Canada 1988.

[5] H. S. M. Coxeter, *Self-dual configurations and regular graphs*, Bull. Amer. Math. Soc., **56** (1950), 413–455.

[6] I. Dejter, *On a* $\{K_4, K_{2,2,2}\}$*-ultrahomogeneous graph*, Australasian Journal of Combinatorics, **44** (2009), 63–76.

[7] I. Dejter, *From the Coxeter graph to the Klein graph*, Journal of Graph Theory, Wyley Online Library.

[8] A. Gardiner, *Homogeneous graphs*, J. Combinatorial Theory (B), **20** (1976), 94–102.

[9] C. Godsil and G. Royle, Algebraic Graph Theory, Springer–Verlag, 2001.

[10] Ja. Ju. Gol'fand and M. H. Klin, *On k-homogeneous graphs*, Algorithmic studies in combinatorics (Russian), **186** (1978), 76–85.

[11] J. L. Gross and J. Yellen eds., Handbook of Graph Theory, CRC Press, 2003.

[12] D. C. Isaksen, C. Jankowski and S. Proctor, *On* K_**-ultrahomogeneous graphs*, Ars Combinatoria, Volume LXXXII, (2007), 83–96.

[13] J. Lauri and R. Scapellato, Topics in graph automorphisms and reconstruction, London Math. Soc., No. 54, Cambridge Univ. Press, 2003.

[14] C. Ronse, *On homogeneous graphs*, J. London Math. Soc., **s2-17** (1978), 375–379.

[15] J. Sheehan, *Smoothly embeddable subgraphs*, J. London Math. Soc., **s2-9** (1974), 212–218.